

ACR/EULAR (2010) Classification Criteria

ACR/EULAR (2009) Classification Criteria for Rheumatoid Arthritis

| <u>Symptom Duration (as reported by patient)</u> | <u>Points</u> |
|--|---------------|
|--|---------------|

- | | |
|-------------|---|
| ▪ < 6 weeks | 0 |
| ▪ > 6 weeks | 1 |

| <u>Joint Distribution</u> | <u>Points</u> |
|---------------------------|---------------|
|---------------------------|---------------|

- | | |
|---|---|
| ▪ 1 large joint | 0 |
| ▪ 2-10 large joints | 1 |
| ▪ 1-3 small joints (with or without involvement of large joints) | 2 |
| ▪ 4-10 small joints (with or without involvement of large joints) | 4 |
| ▪ > 10 joints (at least 1 small joint) | 5 |

| <u>Serology</u> | <u>Points</u> |
|-----------------|---------------|
|-----------------|---------------|

- | | |
|--------------------|---|
| ▪ RF- and CCP- | 0 |
| ▪ Low RF+ or CCP+ | 2 |
| ▪ High RF+ or CCP+ | 3 |

| <u>Acute Phase Reactants</u> | <u>Points</u> |
|------------------------------|---------------|
|------------------------------|---------------|

- | | |
|-----------------------|---|
| ▪ Normal ESR or CRP | 0 |
| ▪ Abnormal ESR or CRP | 1 |

RF: rheumatoid factor. CCP: anti-citrullinated citric peptide. ESR: erythrocyte sedimentation rate. CRP: C-reactive protein. Low: < 3 x upper limit of normal (ULN). High: > 3 x ULN

Requirements: patients who have at least 1 swollen joint, and not better explained by another disease to be applied. A score ≥ 6 points is required for classification as definite RA.

Reference: Aletaha et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology / European League Against Rheumatism collaborative initiative. *Ann Rheum Dis* 2010;69:1580-1588.